

Product datasheet for **RC219935**

UBE2C (NM_181803) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: UBE2C (NM_181803) Human Tagged ORF Clone

Tag: Myc-DDK

Symbol: UBE2C

Synonyms: dj447F3.2; UBCH10

Vector: pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

Cell Selection: Neomycin

ORF Nucleotide Sequence: >RC219935 representing NM_181803
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGGCTTCCAAAACCGCAGCCAGCCGCACTAGCGTCGCCGCCCGTAAAGGAGCTGAGCCGAGCG
GGGGCGCCGCCGGGTCCGGTGGGCAAAGGCTACAGCAGGAGCTGATGACCCTCATGAACCCAACATT
GATAGTCCT

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC219935 representing NM_181803
Red=Cloning site Green=Tags(s)

MASQNRDPAATSVAAARKGAEPSSGAARGPVGKRLQQELMTLMNPTLIVP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-Mlul




[View online »](#)

Cloning Scheme:



Plasmid Map:



ACCN: NM_181803
 ORF Size: 150 bp

| | |
|-------------------------------|---|
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| Components: | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water). |
| Reconstitution Method: | <ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | NM_181803.1 , NP_861519.1 |
| RefSeq Size: | 520 bp |
| RefSeq ORF: | 152 bp |
| Locus ID: | 11065 |
| Cytogenetics: | 20q13.12 |
| Protein Families: | Druggable Genome |
| Protein Pathways: | Ubiquitin mediated proteolysis |
| MW: | 4.9 kDa |
| Gene Summary: | The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, ubiquitin-conjugating enzymes, and ubiquitin-protein ligases. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. The encoded protein is required for the destruction of mitotic cyclins and for cell cycle progression, and may be involved in cancer progression. Multiple transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene have been defined on chromosomes 4, 14, 15, 18, and 19. [provided by RefSeq, Aug 2013] |