

## **Product datasheet for RC219954L2**

# BAX (NM\_004324) Human Tagged Lenti ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** BAX (NM\_004324) Human Tagged Lenti ORF Clone

Tag: mGFP Symbol: BAX

Synonyms: BCL2L4

Mammalian Cell None

Selection:

**Vector:** pLenti-C-mGFP (PS100071)

E. coli Selection: Chloramphenicol (34 ug/mL)


ORF Nucleotide The ORF insert of this clone is exactly the same as(RC219954).

Sequence:

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF.

**ACCN:** NM\_004324

ORF Size: 654 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

### BAX (NM\_004324) Human Tagged Lenti ORF Clone - RC219954L2

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 004324.3</u>

RefSeq Size: 891 bp
RefSeq ORF: 657 bp
Locus ID: 581

 UniProt ID:
 Q07812

 Cytogenetics:
 19q13.33

**Domains:** Bcl-2

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Amyotrophic lateral sclerosis (ALS), Apoptosis, Colorectal cancer, Huntington's disease,

Neurotrophin signaling pathway, p53 signaling pathway, Pathways in cancer, Prion diseases

MW: 24 kDa

**Gene Summary:** The protein encoded by this gene belongs to the BCL2 protein family. BCL2 family members

a wide variety of cellular activities. This protein forms a heterodimer with BCL2, and functions as an apoptotic activator. The association and the ratio of BAX to BCL2 also determines survival or death of a cell following an apoptotic stimulus. This protein is reported to interact with, and increase the opening of, the mitochondrial voltage-dependent anion channel


form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in

expression of this gene is regulated by the tumor suppressor P53 and has been shown to be involved in P53-mediated apoptosis. Multiple alternatively spliced transcript variants, which encode different isoforms, have been reported for this gene. [provided by RefSeq, Dec 2019]

(VDAC), which leads to the loss in membrane potential and the release of cytochrome c. The



## **Product images:**



Circular map for RC219954L2