

## Product datasheet for RC220025L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## EIF4G1 (NM\_182917) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: EIF4G1 (NM 182917) Human Tagged ORF Clone Lentiviral Particle

Symbol: EIF4G1

Synonyms: EIF-4G1; EIF4F; EIF4G; EIF4G1; P220; PARK18

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_182917 **ORF Size:** 4797 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC220025).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 182917.2</u>

 RefSeq Size:
 5539 bp

 RefSeq ORF:
 4803 bp

 Locus ID:
 1981

 UniProt ID:
 Q04637

 Cytogenetics:
 3q27.1

**Protein Pathways:** Viral myocarditis

**MW:** 175.3 kDa







## **Gene Summary:**

The protein encoded by this gene is a component of the multi-subunit protein complex EIF4F. This complex facilitates the recruitment of mRNA to the ribosome, which is a rate-limiting step during the initiation phase of protein synthesis. The recognition of the mRNA cap and the ATP-dependent unwinding of 5'-terminal secondary structure is catalyzed by factors in this complex. The subunit encoded by this gene is a large scaffolding protein that contains binding sites for other members of the EIF4F complex. A domain at its N-terminus can also interact with the poly(A)-binding protein, which may mediate the circularization of mRNA during translation. Alternative splicing results in multiple transcript variants, some of which are derived from alternative promoter usage. [provided by RefSeq, Aug 2010]