

### Product datasheet for RC220060L1V

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Transferrin Receptor 2 (TFR2) (NM\_003227) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** Transferrin Receptor 2 (TFR2) (NM\_003227) Human Tagged ORF Clone Lentiviral Particle

**Symbol:** Transferrin Receptor 2

Synonyms: HFE3; TFRC2

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM\_003227

ORF Size: 2403 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC220060).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally accurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 003227.3

 RefSeq Size:
 2891 bp

 RefSeq ORF:
 2406 bp

 Locus ID:
 7036

 UniProt ID:
 Q9UP52

 Cytogenetics:
 7q22.1

**Protein Families:** Druggable Genome, Protease, Transmembrane

MW: 88.6 kDa





# Transferrin Receptor 2 (TFR2) (NM\_003227) Human Tagged ORF Clone Lentiviral Particle – RC220060L1V

#### **Gene Summary:**

This gene encodes a single-pass type II membrane protein, which is a member of the transferrin receptor-like family. This protein mediates cellular uptake of transferrin-bound iron, and may be involved in iron metabolism, hepatocyte function and erythrocyte differentiation. Mutations in this gene have been associated with hereditary hemochromatosis type III. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, May 2011]