

Product datasheet for RC220061L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MTMR3 (NM_153051) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: MTMR3 (NM_153051) Human Tagged ORF Clone Lentiviral Particle

Symbol: MTMR3

Synonyms: FYVE-DSP1; ZFYVE10

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_153051 **ORF Size:** 3483 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC220061).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 153051.2

 RefSeq Size:
 8897 bp

 RefSeq ORF:
 3486 bp

 Locus ID:
 8897

 UniProt ID:
 Q13615

 Cytogenetics:
 22q12.2

Domains: FYVE, PTPc_motif

Protein Families: Druggable Genome, Phosphatase





ORIGENE

MW: 129.4 kDa

Gene Summary:

This gene encodes a member of the myotubularin dual specificity protein phosphatase gene family. The encoded protein is structurally similar to myotubularin but in addition contains a FYVE domain and an N-terminal PH-GRAM domain. The protein can self-associate and also form heteromers with another myotubularin related protein. The protein binds to phosphoinositide lipids through the PH-GRAM domain, and can hydrolyze phosphatidylinositol(3)-phosphate and phosphatidylinositol(3,5)-biphosphate in vitro. The encoded protein has been observed to have a perinuclear, possibly membrane-bound, distribution in cells, but it has also been found free in the cytoplasm. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]