

Product datasheet for RC220079L1

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Wilms Tumor Protein (WT1) (NM_000378) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Wilms Tumor Protein (WT1) (NM_000378) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: Wilms Tumor Protein

Synonyms: AWT1; GUD; NPHS4; WAGR; WIT-2; WT33

Mammalian Cell None

Selection:

Vector:pLenti-C-Myc-DDK (PS100064)E. coli Selection:Chloramphenicol (34 ug/mL)


ORF Nucleotide The ORF insert of this clone is exactly the same as(RC220079).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_000378

ORF Size: 1491 bp



Wilms Tumor Protein (WT1) (NM_000378) Human Tagged Lenti ORF Clone - RC220079L1

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 000378.3</u>

 RefSeq Size:
 2977 bp

 RefSeq ORF:
 1509 bp

 Locus ID:
 7490

 UniProt ID:
 P19544

Cytogenetics: 11p13

Protein Families: Druggable Genome, Transcription Factors

MW: 54.3 kDa


Gene Summary: This gene encodes a transcription factor that contains four zinc-finger motifs at the C-

terminus and a proline/glutamine-rich DNA-binding domain at the N-terminus. It has an essential role in the normal development of the urogenital system, and it is mutated in a small subset of patients with Wilms tumor. This gene exhibits complex tissue-specific and polymorphic imprinting pattern, with biallelic, and monoallelic expression from the maternal and paternal alleles in different tissues. Multiple transcript variants have been described. In several variants, there is evidence for the use of a non-AUG (CUG) translation initiation codon upstream of, and in-frame with the first AUG. Authors of PMID:7926762 also provide evidence that WT1 mRNA undergoes RNA editing in human and rat, and that this process is tissue-


restricted and developmentally regulated. [provided by RefSeq, Mar 2015]



Product images:



Circular map for RC220079L1



Double digestion of RC220079L1 using Sgfl and Mlul $\,$