

Product datasheet for RC220100L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

LY108 (SLAMF6) (NM_052931) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: LY108 (SLAMF6) (NM_052931) Human Tagged ORF Clone Lentiviral Particle

Symbol: LY108

Synonyms: CD352; KALI; KALIb; Ly108; NTB-A; NTBA; SF2000

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 052931

ORF Size: 993 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC220100).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 052931.3

 RefSeq Size:
 2776 bp

 RefSeq ORF:
 996 bp

 Locus ID:
 114836

 UniProt ID:
 Q96DU3

Cytogenetics: 1q23.2-q23.3

Domains: ig, IG

Protein Families: Druggable Genome, Transmembrane





MW: 34.2 kDa

Gene Summary:

The protein encoded by this gene is a type I transmembrane protein, belonging to the CD2 subfamily of the immunoglobulin superfamily. This encoded protein is expressed on Natural killer (NK), T, and B lymphocytes. It undergoes tyrosine phosphorylation and associates with the Src homology 2 domain-containing protein (SH2D1A) as well as with SH2 domain-containing phosphatases (SHPs). It functions as a coreceptor in the process of NK cell activation. It can also mediate inhibitory signals in NK cells from X-linked lymphoproliferative patients. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, May 2010]