

## Product datasheet for RC220107L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PBR (TSPO) (NM\_000714) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: PBR (TSPO) (NM\_000714) Human Tagged ORF Clone Lentiviral Particle

Symbol: PBR

Synonyms: BPBS; BZRP; DBI; IBP; MBR; mDRC; PBR; PBS; pk18; PKBS; PTBR

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM\_000714

ORF Size: 507 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC220107).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 000714.4

 RefSeq Size:
 921 bp

 RefSeq ORF:
 510 bp

 Locus ID:
 706

 UniProt ID:
 P30536

 Cytogenetics:
 22q13.2

**Domains:** TspO\_MBR

**Protein Families:** Druggable Genome, Transmembrane





## PBR (TSPO) (NM\_000714) Human Tagged ORF Clone Lentiviral Particle - RC220107L2V

**Protein Pathways:** Neuroactive ligand-receptor interaction

**MW:** 18.8 kDa

**Gene Summary:** Present mainly in the mitochondrial compartment of peripheral tissues, the protein encoded

by this gene interacts with some benzodiazepines and has different affinities than its endogenous counterpart. The protein is a key factor in the flow of cholesterol into mitochondria to permit the initiation of steroid hormone synthesis. Alternatively spliced transcript variants have been reported; one of the variants lacks an internal exon and is considered non-coding, and the other variants encode the same protein. [provided by

RefSeq, Feb 2012]