

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC220122L4V

## BAG2 (NM\_004282) Human Tagged ORF Clone Lentiviral Particle

## Product data:

Product Type:	Lentiviral Particles
Product Name:	BAG2 (NM_004282) Human Tagged ORF Clone Lentiviral Particle
Symbol:	BAG2
Synonyms:	BAG-2; dJ417l1.2
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_004282
ORF Size:	633 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC220122).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 004282.2</u>
RefSeq Size:	2068 bp
RefSeq ORF:	636 bp
Locus ID:	9532
UniProt ID:	<u>O95816</u>
Cytogenetics:	6p12.1
Protein Families:	Druggable Genome
MW:	23.8 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary: BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The predicted BAG2 protein contains 211 amino acids. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US