

## Product datasheet for RC220233L4

### TIAM1 (NM\_003253) Human Tagged Lenti ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	TIAM1 (NM_003253) Human Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	TIAM1
Synonyms:	TIAM-1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC220233).
Restriction Sites:	SgfI-RsrII
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.

ACCN:	NM_003253
ORF Size:	4773 bp




[View online »](#)

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_003253.1</a>
<b>RefSeq Size:</b>	5521 bp
<b>RefSeq ORF:</b>	4776 bp
<b>Locus ID:</b>	7074
<b>UniProt ID:</b>	<a href="#">Q13009</a>
<b>Cytogenetics:</b>	21q22.11
<b>Domains:</b>	RhoGEF, PDZ, PH, RBD
<b>Protein Families:</b>	Druggable Genome
<b>Protein Pathways:</b>	Chemokine signaling pathway, Regulation of actin cytoskeleton
<b>MW:</b>	177.3 kDa

**Gene Summary:**

This gene encodes a RAC1-specific guanine nucleotide exchange factor (GEF). GEFs mediate the exchange of guanosine diphosphate (GDP) for guanosine triphosphate (GTP). The binding of GTP induces a conformational change in RAC1 that allows downstream effectors to bind and transduce a signal. This gene thus regulates RAC1 signaling pathways that affect cell shape, migration, adhesion, growth, survival, and polarity, as well as influencing actin cytoskeletal formation, endocytosis, and membrane trafficking. This gene thus plays an important role in cell invasion, metastasis, and carcinogenesis. In addition to RAC1, the encoded protein activates additional Rho-like GTPases such as CDC42, RAC2, RAC3 and RHOA. This gene encodes multiple protein isoforms that experience a diverse array of intramolecular, protein-protein, and phosphorylation interactions as well as phosphoinositide binding. Both the longer and shorter isoforms have C-terminal Dbl homology (DH) and pleckstrin homology (PH) domains while only the longer isoforms of this gene have the N-terminal myristoylation site and the downstream N-terminal PH domain, ras-binding domain (RBD), and PSD-95/DlgA/ZO-1 (PDZ) domain. [provided by RefSeq, Jul 2017]

**Product images:**


Circular map for RC220233L4