

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC220248L4V

Dectin 1 (CLEC7A) (NM_197950) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Dectin 1 (CLEC7A) (NM_197950) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Dectin 1
Synonyms:	BGR; CANDF4; CD369; CLECSF12; DECTIN1; SCARE2
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_197950
ORF Size:	504 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC220248).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 197950.1</u>
RefSeq Size:	2355 bp
RefSeq ORF:	507 bp
Locus ID:	64581
UniProt ID:	<u>Q9BXN2</u>
Cytogenetics:	12p13.2
Protein Families:	Druggable Genome, Transmembrane
MW:	19 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

CRIGENE Dectin 1 (CLEC7A) (NM_197950) Human Tagged ORF Clone Lentiviral Particle – RC220248L4V

Gene Summary:This gene encodes a member of the C-type lectin/C-type lectin-like domain (CTL/CTLD)
superfamily. The encoded glycoprotein is a small type II membrane receptor with an
extracellular C-type lectin-like domain fold and a cytoplasmic domain with an
immunoreceptor tyrosine-based activation motif. It functions as a pattern-recognition
receptor that recognizes a variety of beta-1,3-linked and beta-1,6-linked glucans from fungi
and plants, and in this way plays a role in innate immune response. Alternate transcriptional
splice variants, encoding different isoforms, have been characterized. This gene is closely
linked to other CTL/CTLD superfamily members on chromosome 12p13 in the natural killer
gene complex region. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US