

## Product datasheet for **RC220309L3V**

### PLA2G2F (NM\_022819) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	PLA2G2F (NM_022819) Human Tagged ORF Clone Lentiviral Particle
Symbol:	PLA2G2F
Synonyms:	GIIFsPLA2; sPLA2-IIF
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_022819
ORF Size:	633 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC220309).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_022819.3</a>
RefSeq Size:	2737 bp
RefSeq ORF:	636 bp
Locus ID:	64600
UniProt ID:	<a href="#">Q9BZM2</a>
Cytogenetics:	1p36.12
Protein Families:	Druggable Genome, Secreted Protein



[View online »](#)

<b>Protein Pathways:</b>	alpha-Linolenic acid metabolism, Arachidonic acid metabolism, Ether lipid metabolism, Fc epsilon RI signaling pathway, Glycerophospholipid metabolism, GnRH signaling pathway, Linoleic acid metabolism, Long-term depression, MAPK signaling pathway, Metabolic pathways, Vascular smooth muscle contraction, VEGF signaling pathway
<b>MW:</b>	23.1 kDa
<b>Gene Summary:</b>	May play a role in lipid mediator production in inflammatory conditions, by providing arachidonic acid to downstream cyclooxygenases and lipoxygenases (By similarity). Phospholipase A2, which catalyzes the calcium-dependent hydrolysis of the 2-acyl groups in 3-sn-phosphoglycerides (PubMed:11112443). Hydrolyzes phosphatidylethanolamine more efficiently than phosphatidylcholine, with only a modest preference for arachidonic acid versus linoelic acid at the sn-2 position. Comparable activity toward 1-palmitoyl-2-oleoyl-phosphatidylserine vesicles to that toward 1-palmitoyl-2-oleoyl-phosphatidylglycerol (By similarity). Hydrolyzes phosphatidylglycerol versus phosphatidylcholine with a 15-fold preference (PubMed:11112443).[UniProtKB/Swiss-Prot Function]