

Product datasheet for RC220323L2

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

C18orf1 (LDLRAD4) (NM_004338) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: C18orf1 (LDLRAD4) (NM_004338) Human Tagged Lenti ORF Clone

Tag:mGFPSymbol:C18orf1Synonyms:C18orf1

Mammalian Cell None

Selection:

Vector: pLenti-C-mGFP (PS100071)


E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC220323).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:






^{*} The last codon before the Stop codon of the ORF.





Plasmid Map:



ACCN: NM_004338

ORF Size: 744 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore,

OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

NM 004338.3, NP 004329.1 RefSeq:

RefSeq Size: 8086 bp RefSeq ORF: 746 bp Locus ID: 753

Cytogenetics: 18p11.21

Protein Families: Druggable Genome, Transmembrane

MW: 27.6 kDa

Gene Summary: Functions as a negative regulator of TGF-beta signaling and thereby probably plays a role in

> cell proliferation, differentiation, apoptosis, motility, extracellular matrix production and immunosuppression. In the canonical TGF-beta pathway, ZFYVE9/SARA recruits the

intracellular signal transducer and transcriptional modulators SMAD2 and SMAD3 to the TGFbeta receptor. Phosphorylated by the receptor, SMAD2 and SMAD3 then form a heteromeric complex with SMAD4 that translocates to the nucleus to regulate transcription. Through interaction with SMAD2 and SMAD3, LDLRAD4 may compete with ZFYVE9 and SMAD4 and

prevent propagation of the intracellular signal.[UniProtKB/Swiss-Prot Function]