

Product datasheet for RC220352L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

EPHA8 (NM_020526) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: EPHA8 (NM_020526) Human Tagged ORF Clone Lentiviral Particle

Symbol: EPHA8

Synonyms: EEK; EK3; HEK3

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_020526 **ORF Size:** 3015 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC220352).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 020526.3

 RefSeq Size:
 4996 bp

 RefSeq ORF:
 3018 bp

 Locus ID:
 2046

 UniProt ID:
 P29322

 Cytogenetics:
 1p36.12

Domains: pkinase, EPH_lbd, TyrKc, SAM, S_TKc, FN3

Protein Families: Druggable Genome, Protein Kinase, Transmembrane







Protein Pathways: Axon guidance

MW: 108.1 kDa

Gene Summary: This gene encodes a member of the ephrin receptor subfamily of the protein-tyrosine kinase

family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. The protein encoded by this gene functions as a receptor for ephrin A2, A3

and A5 and plays a role in short-range contact-mediated axonal guidance during development of the mammalian nervous system. [provided by RefSeq, Jul 2008]