

## **Product datasheet for RC220388**

#### OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

# HP1 gamma (CBX3) (NM\_016587) Human Tagged ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

Product Name: HP1 gamma (CBX3) (NM\_016587) Human Tagged ORF Clone

Tag: Myc-DDK

Symbol: HP1 gamma

Synonyms: HECH; HP1-GAMMA; HP1Hs-gamma

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>RC220388 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC220388 protein sequence

Red=Cloning site Green=Tags(s)

MASNKTTLQKMGKKQNGKSKKVEEAEPEEFVVEKVLDRRVVNGKVEYFLKWKGFTDADNTWEPEENLDCP ELIEAFLNSQKAGKEKDGTKRKSLSDSESDDSKSKKKRDAADKPRGFARGLDPERIIGATDSSGELMFLM

KWKDSDEADLVLAKEANMKCPQIVIAFYEERLTWHSCPEDEAQ

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: <a href="https://cdn.origene.com/chromatograms/mk6299">https://cdn.origene.com/chromatograms/mk6299</a> f08.zip






**Restriction Sites:** 

Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_016587

ORF Size: 549 bp

**OTI Disclaimer:** 

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:customport@origene.com">customport@origene.com</a> or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

**OTI Annotation:** 

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** 

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.



**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

**RefSeq:** <u>NM 016587.4</u>

RefSeq Size: 2102 bp
RefSeq ORF: 552 bp
Locus ID: 11335
UniProt ID: Q13185
Cytogenetics: 7p15.2
Domains: CHROMO

**Protein Families:** Druggable Genome, Transcription Factors


MW: 20.8 kDa

**Gene Summary:** At the nuclear envelope, the nuclear lamina and heterochromatin are adjacent to the inner

nuclear membrane. The protein encoded by this gene binds DNA and is a component of heterochromatin. This protein also can bind lamin B receptor, an integral membrane protein found in the inner nuclear membrane. The dual binding functions of the encoded protein may explain the association of heterochromatin with the inner nuclear membrane. This protein binds histone H3 tails methylated at Lys-9 sites. This protein is also recruited to sites of ultraviolet-induced DNA damage and double-strand breaks. Two transcript variants encoding the same protein but differing in the 5' UTR, have been found for this gene.


[provided by RefSeq, Mar 2011]


### **Product images:**



Circular map for RC220388







Western blot validation of overexpression lysate (Cat# [LY413887]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC220388 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

Coomassie blue staining of purified CBX3 protein (Cat# [TP320388]). The protein was produced from HEK293T cells transfected with CBX3 cDNA clone (Cat# RC220388) using MegaTran 2.0 (Cat# [TT210002]).