

Product datasheet for **RC220400L4V**

LIM kinase 2 (LIMK2) (NM_016733) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	LIM kinase 2 (LIMK2) (NM_016733) Human Tagged ORF Clone Lentiviral Particle
Symbol:	LIM kinase 2
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_016733
ORF Size:	1851 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC220400).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_016733.2
RefSeq Size:	3848 bp
RefSeq ORF:	1854 bp
Locus ID:	3985
UniProt ID:	P53671
Cytogenetics:	22q12.2
Domains:	pkinase, TyrKc, PDZ, LIM, S_TKc
Protein Families:	Druggable Genome, Protein Kinase
Protein Pathways:	Axon guidance, Fc gamma R-mediated phagocytosis, Regulation of actin cytoskeleton



[View online »](#)

MW: 69.7 kDa

Gene Summary: There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. The protein encoded by this gene is phosphorylated and activated by ROCK, a downstream effector of Rho, and the encoded protein, in turn, phosphorylates cofilin, inhibiting its actin-depolymerizing activity. It is thought that this pathway contributes to Rho-induced reorganization of the actin cytoskeleton. At least three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]