

## Product datasheet for RC220453L4

### PDIR (PDIA5) (NM\_006810) Human Tagged Lenti ORF Clone

#### Product data:

|                           |  |
|---------------------------|--|
| Product Type:             | Expression Plasmids  |
| Product Name:             | PDIR (PDIA5) (NM_006810) Human Tagged Lenti ORF Clone          |
| Tag:                      | mGFP   |
| Symbol:                   | PDIR   |
| Synonyms:                 | PDIR   |
| Mammalian Cell Selection: | Puromycin  |
| Vector:                   | pLenti-C-mGFP-P2A-Puro (PS100093)                              |
| E. coli Selection:        | Chloramphenicol (34 ug/mL)                                     |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(RC220453). |
| Restriction Sites:        | SgfI-MluI  |
| Cloning Scheme:           |  |

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.


|           |           |
|-----------|-----------|
| ACCN:     | NM_006810 |
| ORF Size: | 1557 bp   |



[View online »](#)

|                               |   |
|-------------------------------|---|
| <b>OTI Disclaimer:</b>        | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>  |
| <b>OTI Annotation:</b>        | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.  |
| <b>Components:</b>            | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).  |
| <b>Reconstitution Method:</b> | <ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>   |
| <b>RefSeq:</b>                | <a href="#">NM_006810.3</a>   |
| <b>RefSeq Size:</b>           | 1873 bp   |
| <b>RefSeq ORF:</b>            | 1560 bp   |
| <b>Locus ID:</b>              | 10954   |
| <b>UniProt ID:</b>            | <a href="#">Q14554</a>  |
| <b>Cytogenetics:</b>          | 3q21.1  |
| <b>Domains:</b>               | thioered  |
| <b>Protein Families:</b>      | Druggable Genome  |
| <b>MW:</b>                    | 59.6 kDa  |
| <b>Gene Summary:</b>          | This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions. The encoded protein has an N-terminal ER-signal sequence, three catalytically active thioredoxin (TRX) domains, a TRX-like domain, and a C-terminal ER-retention sequence. The N-terminal TRX-like domain is the primary binding site for the major ER chaperone calreticulin and possibly other proteins and substrates as well. Alternative splicing results in multiple protein- and non-protein-coding transcript variants. [provided by RefSeq, Dec 2016] |

Product images:



Circular map for RC220453L4