

#### OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for RC220500L4

## ErbB 4 (ERBB4) (NM\_001042599) Human Tagged Lenti ORF Clone

## **Product data:**

Product Type:	Expression Plasmids
Product Name:	ErbB 4 (ERBB4) (NM_001042599) Human Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	ErbB 4
Synonyms:	ALS19; HER4; p180erbB4
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC220500).
<b>Restriction Sites:</b>	Sgfl-Mlul
Cloning Scheme:	
	Cloning sites used for ORF Shuttling:
	Sgf I         ORF         Mlu I          GCG ATC GCC         ATG// NNN         ACG CGT



ACCN: ORF Size: NM\_001042599 3876 bp



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


	rbB 4 (ERBB4) (NM_001042599) Human Tagged Lenti ORF Clone – RC220500L4
OTI Disclaimer:	Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <u>custsupport@origene.com</u> or by calling 301.340.3188 option 3 for pricing and delivery.
	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Me	<ul> <li>thod: 1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ul>
RefSeq:	<u>NM 001042599.1, NP 001036064.1</u>
RefSeq Size:	11893 bp
RefSeq ORF:	3879 bp
Locus ID:	2066
UniProt ID:	<u>Q15303</u>
Cytogenetics:	2q34
Protein Families:	Druggable Genome, Protein Kinase, Transmembrane
Protein Pathways:	Calcium signaling pathway, Endocytosis, ErbB signaling pathway
MW:	145.2 kDa

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


#### Scheme Content Content

Gene Summary:This gene is a member of the Tyr protein kinase family and the epidermal growth factor<br/>receptor subfamily. It encodes a single-pass type I membrane protein with multiple cysteine<br/>rich domains, a transmembrane domain, a tyrosine kinase domain, a phosphotidylinositol-3<br/>kinase binding site and a PDZ domain binding motif. The protein binds to and is activated by<br/>neuregulins and other factors and induces a variety of cellular responses including<br/>mitogenesis and differentiation. Multiple proteolytic events allow for the release of a<br/>cytoplasmic fragment and an extracellular fragment. Mutations in this gene have been<br/>associated with cancer. Alternatively spliced variants which encode different protein isoforms<br/>have been described; however, not all variants have been fully characterized. [provided by<br/>RefSeq, Jul 2008]

#### **Product images:**



Circular map for RC220500L4



Double digestion of RC220500L4 using Sgfl and Mlul

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US