

Product datasheet for RC220534L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

POLR3H (NM_138338) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: POLR3H (NM 138338) Human Tagged ORF Clone Lentiviral Particle

Symbol: POLR3H

Synonyms: C25; RPC8; RPC22.9

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_138338

ORF Size: 612 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC220534).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 138338.2

 RefSeq Size:
 4186 bp

 RefSeq ORF:
 615 bp

 Locus ID:
 171568

 UniProt ID:
 Q9Y535

 Cytogenetics:
 22q13.2

Domains: RNA_pol_Rpb7_N

Protein Families: Transcription Factors





POLR3H (NM_138338) Human Tagged ORF Clone Lentiviral Particle - RC220534L3V

Protein Pathways: Cytosolic DNA-sensing pathway, Metabolic pathways, Purine metabolism, Pyrimidine

metabolism, RNA polymerase

MW: 22.9 kDa

Gene Summary: DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four

ribonucleoside triphosphates as substrates. Specific peripheric component of RNA

polymerase III which synthesizes small RNAs, such as 5S rRNA and tRNAs. Plays a key role in sensing and limiting infection by intracellular bacteria and DNA viruses. Acts as nuclear and cytosolic DNA sensor involved in innate immune response. Can sense non-self dsDNA that serves as template for transcription into dsRNA. The non-self RNA polymerase III transcripts, such as Epstein-Barr virus-encoded RNAs (EBERS) induce type I interferon and NF- Kappa-B

through the RIG-I pathway (By similarity).[UniProtKB/Swiss-Prot Function]