

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC220544L3V

Cyclin C (CCNC) (NM_005190) Human Tagged ORF Clone Lentiviral Particle

Product data:

| Product Type: | Lentiviral Particles |
|------------------------------|---|
| Product Name: | Cyclin C (CCNC) (NM_005190) Human Tagged ORF Clone Lentiviral Particle |
| Symbol: | Cyclin C |
| Synonyms: | CycC; hSRB11; SRB11 |
| Mammalian Cell Selection: | Puromycin |
| Vector: | pLenti-C-Myc-DDK-P2A-Puro (PS100092) |
| Tag: | Myc-DDK |
| ACCN: | NM_005190 |
| ORF Size: | 849 bp |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(RC220544). |
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u> |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| RefSeq: | <u>NM 005190.3</u> |
| RefSeq Size: | 2321 bp |
| RefSeq ORF: | 852 bp |
| Locus ID: | 892 |
| UniProt ID: | <u>P24863</u> |
| Cytogenetics: | 6q16.2 |
| Domains: | CYCLIN, cyclin |
| Protein Families: | Druggable Genome, Transcription Factors |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

| | Cyclin C (CCNC) (NM_005190) Human Tagged ORF Clone Lentiviral Particle – RC220544L3V |
|---------------|--|
| MW: | 33.1 kDa |
| Gene Summary: | The protein encoded by this gene is a member of the cyclin family of proteins. The encoded protein interacts with cyclin-dependent kinase 8 and induces the phophorylation of the carboxy-terminal domain of the large subunit of RNA polymerase II. The level of mRNAs for this gene peaks in the G1 phase of the cell cycle. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008] |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US