

Product datasheet for **RC220561**

Histone H2A Bbd (H2AFB3) (NM_080720) Human Tagged ORF Clone

Product data:

| | |
|---------------------------|---|
| Product Type: | Expression Plasmids |
| Product Name: | Histone H2A Bbd (H2AFB3) (NM_080720) Human Tagged ORF Clone |
| Tag: | Myc-DDK |
| Symbol: | Histone H2A Bbd |
| Synonyms: | H2AB2; H2ABBD; H2AFB; H2AFB3 |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-Entry (PS100001) |
| E. coli Selection: | Kanamycin (25 ug/mL) |
| ORF Nucleotide Sequence: | >RC220561 representing NM_080720 Red=Cloning site Blue=ORF Green=Tags(s) TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCCGCC CGATCGCC ATGCCGAGGAGGAGACGCCGAGGGTCTCCGGTGTGGCGGCCGGGGCGGACCTGCTCTCGCACCGTCCGAGCGGAGCTTTCGTTTTTCAGTGAGCCAGGTGGAGCGCAGTCTACGGGAGGGCCACTACGCTCAGCGCCTGAGTCGCACGGCGCCGGTCTACCTCGCTGCGGTTATTGAGTACCTGACGGCCAAGTCTGGAGCTGGCGGCAACGAGGCCAGAACAGCGGAGAGCGGAACATCACTCCCCTGCTGCTGGACATGGTGGTTCACAACGACAGGCTACTGAGCACCCCTTTCAACACGACCACCATCTCTCAAGTGCCCTGGCGAGGAC ACGCGT ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA >RC220561 representing NM_080720 Red=Cloning site Green=Tags(s) MPRRRRRRGSSGAGGRGRTCSRTVRAELSFVSQVERSLREGHYAQRSLRTAPVYLAHVIEYLAKVLELAGNEAQNISGERNITPLLLDMVVHNDRLSTLFNTTISQVAPGED TR TRPLEQKLISEEDLAANDILDYKDDDDKV |
| Protein Sequence: | >RC220561 representing NM_080720 Red=Cloning site Green=Tags(s) MPRRRRRRGSSGAGGRGRTCSRTVRAELSFVSQVERSLREGHYAQRSLRTAPVYLAHVIEYLAKVLELAGNEAQNISGERNITPLLLDMVVHNDRLSTLFNTTISQVAPGED TR TRPLEQKLISEEDLAANDILDYKDDDDKV |
| Chromatograms: | https://cdn.origene.com/chromatograms/mk8102_a10.zip |
| Restriction Sites: | SgfI-MluI |




[View online »](#)


UniProt ID: [P0C5Z0](#)
Cytogenetics: Xq28
Protein Pathways: Systemic lupus erythematosus
MW: 12.7 kDa

Gene Summary: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-independent histone that is a member of the histone H2A family. This gene is part of a region that is repeated three times on chromosome X, once in intron 22 of the F8 gene and twice closer to the Xq telomere. This record represents the most telomeric copy. [provided by RefSeq, Oct 2015]

Product images:



Circular map for RC220561



Western blot validation of overexpression lysate (Cat# [LY409042]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC220561 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).