

Product datasheet for RC220561L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Histone H2A Bbd (H2AFB3) (NM 080720) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Histone H2A Bbd (H2AFB3) (NM_080720) Human Tagged ORF Clone Lentiviral Particle

Symbol: Histone H2A Bbd

Synonyms: H2AB2; H2ABBD; H2AFB; H2AFB3

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_080720

ORF Size: 345 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC220561).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 080720.1

RefSeq Size: 539 bp
RefSeq ORF: 348 bp
Locus ID: 83740
UniProt ID: POC5Z0
Cytogenetics: Xq28

Protein Pathways: Systemic lupus erythematosus

MW: 12.7 kDa





Histone H2A Bbd (H2AFB3) (NM_080720) Human Tagged ORF Clone Lentiviral Particle – RC220561L4V

Gene Summary:

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-independent histone that is a member of the histone H2A family. This gene is part of a region that is repeated three times on chromosome X, once in intron 22 of the F8 gene and twice closer to the Xq telomere. This record represents the most telomeric copy. [provided by RefSeq, Oct 2015]