

Product datasheet for **RC220599**

PKC beta 1 (PRKCB) (NM_212535) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	PKC beta 1 (PRKCB) (NM_212535) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	PKC beta 1
Synonyms:	PKC-beta; PKCB; PKCbeta; PKCI(2); PRKCB1; PRKCB2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide
Sequence:

>RC220599 representing NM_212535
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGGCTGACCCGGCTGCGGGGCCCGCCGAGCGAGGGCGAGGAGACACCGTGCCTTCGCCCGCAAAG
GCGCCCTCCGGCAGAAGAACGTGCATGAGGTCAAGAACCACAAATTCACCGCCCGCTTCTCAAGCAGCC
CACCTTCTGCAGCCACTGCACCGACTTCATCTGGGGCTTCGGAAGCAGGGATTCCAGTGCCAAGTTTGC
TGCTTTGTGGTGACAAGCGGTGCCATGAATTTGTACATTCTCCTGCCCTGGCGCTGACAAGGGTCCAG
CCTCCGATGACCCCGCAGCAAACACAAGTTAAGATCCACACGTAATCCAGCCACGTTTTGTGACCA
CTGTGGTCACTGCTGTATGGACTATCCACCAGGGGATGAAATGTGACACCTGCATGATGAATGTGCAC
AAGCGTCCGTGATGAATGTTCCAGCCTGTGTGGCACGGACCACACGGAGCGCCGCGCCGCATCTACA
TCCAGGCCACATCGACAGGGACGTCTCATTGTCTCGTAAGAGATGCTAAAACTTTGTACCTATGGA
CCCAATGGCCTGTCAGATCCCTACGTAAGTAACTGAACTGATTCCCGATCCAAAAGTGAGAGCAAACAG
AAGACCAAAACCATCAAATGCTCCCTCAACCTGAGTGAATGAGACATTTAGATTTAGCTGAAAGAAT
CGGACAAAAGACAGAAGACTGTCAGTAGAGATTTGGGATTTGGGATTTGACCAGCAGGAATGACTTCATGGG
ATCTTTGTCTTTGGGATTTCTGAACCTTCAGAAAGCCAGTGTGATGGCTGGTTTAAAGTTACTGAGCCAG
GAGGAAGGCGAGTACTTCAATGTGCCTGTGCCACCAGAAGGAAGTGAGGCCAATGAAGAAGTGGCCGAGA
AATTTGAGAGGGCCAAGATCAGTCAGGGAACCAAGGTCCCGGAAGAAAAGACGACCAACACTGTCTCAA
ATTTGACAACAATGGCAACAGAGACCGGATGAACTGACCGATTTTAACTTCTAATGGTGTGGGGAAA
GGCAGCTTTGGCAAGGTGATGCTTTGAGAAGCAAGGACAGATGAGCTCTATGCTGTGAAGATCCTGA
AGAAGGACGTTGTGATCCAAGATGATGACGTGGAGTGCCTATGGTGGAGAAGCGGGTGTGGCCCTGCC
CGGAAGCCGCCCTTCTGACCCAGCTCCACTCCTGCTTCCAGACCATGGACCGCCTGTACTTTGTGATG
GAGTACGTGAATGGGGGCGACCTCATGTATCACATCCAGCAAGTCGCGCCGTTCAAGGAGCCCCATGCTG
TATTTTACGCTGCAGAAATTGCCATCGGTCTGTTCTTCTTACAGAGTAAGGGCATCATTTACCGTGACCT
AAAAGTGGCAACGTGATGCTCGATTCTGAGGGACACATCAAGATTGCCGATTTTGGCATGTGTAAGGAA
AACATCTGGGATGGGGTGACAACCAAGACATTCTGTGGCACTCCAGACTACATCGCCCCGAGATAATTG
CTTATCAGCCCTATGGGAAGTCCGTGGATTGGTGGGCATTTGGAGTCTGCTGTATGAAATGTTGGCTGG
GCAGGCACCCCTTGAAGGGGAGGATGAAGATGAACTTCCAATCCATCATGGAACACAACGTAGCCTAT
CCCAAGTCTATGTCCAAGGAAGCTGTGGCCATCTGCAAAGGGCTGATGACCAACACCCAGGCAAACGTC
TGGGTTGTGGACCTGAAGGCGAACGTGATATCAAAGAGCATGCATTTTCCGGTATATTGATTGGGAGAA
ACTTGAACGCAAAGAGATCCAGCCCCCTATAAGCCAAAAGCTAGAGACAAGAGAGACACCTCCAATTC
GACAAAGAGTTACACAGACAGCCTGTGGAAGTACCCCACTGATAAACTCTTATCATGAAGTTGGACC
AAAATGAATTTGCTGGCTTCTTATACTAACCAGAGTTTGTCAATATGTG

ACGCGTACGCGGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC220599 representing NM_212535
Red=Cloning site Green=Tags(s)

```
MADPAAGPPPSEGEESTVRFARKGALRQKNVHEVKNHKFTARFFKQPTFCSHCTDFIWGFQKQGFQCQVC
CFVVHKRCHEFVTFSCPGADKGPASDDPRSKHKFKIHTYSPTFCDHCGSLLYGLIHQGMKCDTCMMNVH
KRCVMNVPSLCGTDHTERRGRIYIQAHDIRDVLIVLVRDAKNLVPMDPNGLSDPYVKLKLIPDPKSESKQ
KTKTIKCSLNPEWNETFRFQLKESDKDRRLSVEIWDWDLTSRNDFMGSLSFSGISELQKASVDGWFKLLSQ
EEGEYFNVPVPEGSEANEELRQKFERAKISQGTKVPEEKTNTNTVSKFDNNGNRDRMKLTDNFNLMVLGK
GSFGKVMLSERKGTDELYAVKILKDVVIQDDDVECTMVEKRVLALPGKPPFLTQLHSCFQTMDRLYFVM
EYVNGDLMYHIQQVGRFKEPHAVFYAAEIAIGLFFLQSKGIIYRDLKLDNVMLDSEGHIKIADFGMCKE
NIWDGVTTKTFCGTPDYIAPEIIAYQPYGKSVDDWAFGVLLYEMLAGQAPFEGEDELEFQSIMEHNVAY
PKSMSKEAVAICKGLMTKHPGKRLGCGPEGERDIKEHAFFRYIDWEKLERKEIQPPYKPKARDKRDTSNF
DKEFTRQPVELTPTDKLFIMNLDQNEFAGFSYTNPEFVINV
```


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mg4220_f01.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_212535


ORF Size: 2013 bp

OTI Disclaimer:	Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.
	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_212535.3
RefSeq Size:	2319 bp
RefSeq ORF:	2016 bp
Locus ID:	5579
UniProt ID:	P05771
Cytogenetics:	16p12.2-p12.1
Protein Families:	Druggable Genome, Protein Kinase
Protein Pathways:	B cell receptor signaling pathway, Calcium signaling pathway, Chemokine signaling pathway, ErbB signaling pathway, Fc epsilon RI signaling pathway, Fc gamma R-mediated phagocytosis, Focal adhesion, Gap junction, Glioma, GnRH signaling pathway, Leukocyte transendothelial migration, Long-term depression, Long-term potentiation, MAPK signaling pathway, Melanogenesis, Natural killer cell mediated cytotoxicity, Non-small cell lung cancer, Pathways in cancer, Phosphatidylinositol signaling system, Tight junction, Vascular smooth muscle contraction, VEGF signaling pathway, Vibrio cholerae infection, Wnt signaling pathway
MW:	76.7 kDa


Gene Summary:

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This protein kinase has been reported to be involved in many different cellular functions, such as B cell activation, apoptosis induction, endothelial cell proliferation, and intestinal sugar absorption. Studies in mice also suggest that this kinase may also regulate neuronal functions and correlate fear-induced conflict behavior after stress. Alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]


Product images:



Circular map for RC220599



Western blot validation of overexpression lysate (Cat# [LY403957]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC220599 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified PRKCB protein (Cat# [TP320599]). The protein was produced from HEK293T cells transfected with PRKCB cDNA clone (Cat# RC220599) using MegaTran 2.0 (Cat# [TT210002]).