

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC220664L1V

LCLAT1 (NM_182551) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	LCLAT1 (NM_182551) Human Tagged ORF Clone Lentiviral Particle
Symbol:	LCLAT1
Synonyms:	1AGPAT8; AGPAT8; ALCAT1; HSRG1849; LYCAT; UNQ1849
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_182551
ORF Size:	1242 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC220664).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 182551.3</u>
RefSeq Size:	5077 bp
RefSeq ORF:	1245 bp
Locus ID:	253558
UniProt ID:	Q6UWP7
Cytogenetics:	2p23.1
Protein Families:	Transmembrane



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US Protein Pathways:Ether lipid metabolism, Glycerolipid metabolism, Glycerophospholipid metabolism, Limonene
and pinene degradation, Metabolic pathways, Phenylalanine metabolism, Tyrosine
metabolism

48.7 kDa

MW:

Gene Summary:Exhibits acyl-CoA:lysocardiolipin acyltransferase (ALCAT) activity; catalyzes the reacylation of
lyso-cardiolipin to cardiolipin (CL), a key step in CL remodeling (By similarity). Recognizes both
monolysocardiolipin and dilysocardiolipin as substrates with a preference for linoleoyl-CoA
and oleoyl-CoA as acyl donors (By similarity). Also exhibits 1-acyl-sn-glycerol-3-phosphate
acyltransferase activity (AGPAT) activity; converts 1-acyl-sn-glycerol-3- phosphate
(lysophosphatidic acid or LPA) into 1,2-diacyl-sn-glycerol-3- phosphate (phosphatidic acid or
PA) by incorporating an acyl moiety at the sn-2 position of the glycerol backbone
(PubMed:16620771). Possesses both lysophosphatidylinositol acyltransferase (LPIAT) and
lysophosphatidylglycerol acyltransferase (LPGAT) activities (PubMed:19075029). Required for
establishment of the hematopoietic and endothelial lineages (By similarity).[UniProtKB/Swiss-
Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US