

Product datasheet for RC220664L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

LCLAT1 (NM_182551) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: LCLAT1 (NM_182551) Human Tagged ORF Clone Lentiviral Particle

Symbol: LCLAT1

Synonyms: 1AGPAT8; AGPAT8; ALCAT1; HSRG1849; LYCAT; UNQ1849

Mammalian Cell

Selection:

Puromycin

Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_182551 **ORF Size:** 1242 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC220664).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 182551.3

RefSeq Size: 5077 bp
RefSeq ORF: 1245 bp
Locus ID: 253558
UniProt ID: Q6UWP7
Cytogenetics: 2p23.1

Protein Families: Transmembrane





LCLAT1 (NM_182551) Human Tagged ORF Clone Lentiviral Particle - RC220664L4V

Protein Pathways: Ether lipid metabolism, Glycerolipid metabolism, Glycerophospholipid metabolism, Limonene

and pinene degradation, Metabolic pathways, Phenylalanine metabolism, Tyrosine

metabolism

MW: 48.7 kDa

Gene Summary: Exhibits acyl-CoA:lysocardiolipin acyltransferase (ALCAT) activity; catalyzes the reacylation of

lyso-cardiolipin to cardiolipin (CL), a key step in CL remodeling (By similarity). Recognizes both monolysocardiolipin and dilysocardiolipin as substrates with a preference for linoleoyl-CoA and oleoyl-CoA as acyl donors (By similarity). Also exhibits 1-acyl-sn-glycerol-3-phosphate

acyltransferase activity (AGPAT) activity; converts 1-acyl-sn-glycerol-3- phosphate

(lysophosphatidic acid or LPA) into 1,2-diacyl-sn-glycerol-3- phosphate (phosphatidic acid or

PA) by incorporating an acyl moiety at the sn-2 position of the glycerol backbone

(PubMed:16620771). Possesses both lysophosphatidylinositol acyltransferase (LPIAT) and lysophosphatidylglycerol acyltransferase (LPGAT) activities (PubMed:19075029). Required for establishment of the hematopoietic and endothelial lineages (By similarity).[UniProtKB/Swiss-

Prot Function]