

Product datasheet for RC220692L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Leucyl cystinyl aminopeptidase (LNPEP) (NM_005575) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Leucyl cystinyl aminopeptidase (LNPEP) (NM_005575) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Leucyl cystinyl aminopeptidase

Synonyms: CAP; IRAP; P-LAP; PLAP

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_005575 **ORF Size:** 3075 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC220692).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 005575.2</u>

 RefSeq Size:
 4470 bp

 RefSeq ORF:
 3078 bp

 Locus ID:
 4012

 UniProt ID:
 Q9UIQ6

 Cytogenetics:
 5q15

Domains: Peptidase_M1





Leucyl cystinyl aminopeptidase (LNPEP) (NM_005575) Human Tagged ORF Clone Lentiviral Particle – RC220692L4V

Protein Families: Druggable Genome, Protease, Secreted Protein, Transmembrane

Protein Pathways: Renin-angiotensin system

MW: 117.2 kDa

Gene Summary: This gene encodes a zinc-dependent aminopeptidase that cleaves vasopressin, oxytocin, lys-

bradykinin, met-enkephalin, dynorphin A and other peptide hormones. The protein can be secreted in maternal serum, reside in intracellular vesicles with the insulin-responsive glucose transporter GLUT4, or form a type II integral membrane glycoprotein. The protein catalyzes the final step in the conversion of angiotensinogen to angiotensin IV (AT4) and is also a receptor for AT4. Alternative splicing results in multiple transcript variants encoding different

isoforms. [provided by RefSeq, Jul 2008]