

Product datasheet for **RC220733L4V**

COASY (NM_025233) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	COASY (NM_025233) Human Tagged ORF Clone Lentiviral Particle
Symbol:	COASY
Synonyms:	DPCK; NBIA6; NBP; PCH12; pOV-2; PPAT; UKR1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_025233
ORF Size:	1692 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC220733).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_025233.4 , NP_079509.4
RefSeq Size:	2470 bp
RefSeq ORF:	1695 bp
Locus ID:	80347
UniProt ID:	Q13057
Cytogenetics:	17q21.2
Domains:	CoaE, CTP_transf_2
Protein Pathways:	Metabolic pathways, Pantothenate and CoA biosynthesis



[View online »](#)

MW: 62.1 kDa

Gene Summary: Coenzyme A (CoA) functions as a carrier of acetyl and acyl groups in cells and thus plays an important role in numerous synthetic and degradative metabolic pathways in all organisms. In eukaryotes, CoA and its derivatives are also involved in membrane trafficking and signal transduction. This gene encodes the bifunctional protein coenzyme A synthase (CoAsy) which carries out the last two steps in the biosynthesis of CoA from pantothenic acid (vitamin B5). The phosphopantetheine adenylyltransferase domain of this bifunctional protein catalyzes the conversion of 4'-phosphopantetheine into dephospho-coenzyme A (dpCoA) while its dephospho-CoA kinase domain completes the final step by phosphorylating dpCoA to form CoA. Mutations in this gene are associated with neurodegeneration with brain iron accumulation (NBIA). Alternative splicing results in multiple isoforms. [provided by RefSeq, Apr 2014]