

## Product datasheet for RC220836L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CYP8B1 (NM 004391) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type: Lentiviral Particles** 

**Product Name:** CYP8B1 (NM\_004391) Human Tagged ORF Clone Lentiviral Particle

Symbol:

CP8B: CYP12 Synonyms:

**Mammalian Cell** 

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

mGFP Tag:

NM 004391 ACCN: **ORF Size:** 1503 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC220836).

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of OTI Disclaimer: reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 004391.1

RefSeq Size: 3951 bp RefSeq ORF: 1506 bp Locus ID: 1582

Q9UNU6 **UniProt ID:** Cytogenetics: 3p22.1 **Domains:** 

**Protein Families:** Druggable Genome, P450, Transmembrane

p450



## CYP8B1 (NM\_004391) Human Tagged ORF Clone Lentiviral Particle - RC220836L2V

**Protein Pathways:** Metabolic pathways, PPAR signaling pathway, Primary bile acid biosynthesis

**MW:** 57.9 kDa

**Gene Summary:** This gene encodes a member of the cytochrome P450 superfamily of enzymes. The

cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This endoplasmic reticulum membrane protein catalyzes the conversion of 7 alpha-hydroxy-4-cholesten-3-one into 7-alpha,12-alpha-dihydroxy-4-cholesten-3-one. The balance between these two steroids determines the relative amounts of cholic acid and chenodeoxycholic acid both of which are secreted in the bile and affect the solubility of cholesterol. This gene is unique among the

cytochrome P450 genes in that it is intronless. [provided by RefSeq, Jul 2008]