

Product datasheet for RC220847L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SOCS1 (NM 003745) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: SOCS1 (NM_003745) Human Tagged ORF Clone Lentiviral Particle

Symbol:

AISIMD; CIS1; CISH1; JAB; SOCS-1; SSI-1; SSI1; TIP-3; TIP3 Synonyms:

Mammalian Cell

Selection:

ACCN:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag: NM 003745

ORF Size: 633 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC220847).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 003745.1, NP 003736.1

RefSeq Size: 1216 bp RefSeq ORF: 636 bp Locus ID: 8651 **UniProt ID:** 015524

Cytogenetics: 16p13.13

Domains: SH2, SOCS





SOCS1 (NM_003745) Human Tagged ORF Clone Lentiviral Particle - RC220847L1V

Protein Families: Druggable Genome, Stem cell - Pluripotency, Stem cell relevant signaling - JAK/STAT signaling

pathway

Protein Pathways: Insulin signaling pathway, Jak-STAT signaling pathway, Type II diabetes mellitus, Ubiquitin

mediated proteolysis

MW: 23.4 kDa

Gene Summary: This gene encodes a member of the STAT-induced STAT inhibitor (SSI), also known as

suppressor of cytokine signaling (SOCS), family. SSI family members are cytokine-inducible negative regulators of cytokine signaling. The expression of this gene can be induced by a subset of cytokines, including IL2, IL3 erythropoietin (EPO), CSF2/GM-CSF, and interferon (IFN)-gamma. The protein encoded by this gene functions downstream of cytokine receptors, and takes part in a negative feedback loop to attenuate cytokine signaling. Knockout studies in mice suggested the role of this gene as a modulator of IFN-gamma action, which is required

for normal postnatal growth and survival. [provided by RefSeq, Jul 2008]