

## Product datasheet for **RC220876L3V**

### UCK (UCK1) (NM\_031432) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

|                           |  |
|---------------------------|--|
| Product Type:             | Lentiviral Particles   |
| Product Name:             | UCK (UCK1) (NM_031432) Human Tagged ORF Clone Lentiviral Particle  |
| Symbol:                   | UCK  |
| Synonyms:                 | URK1   |
| Mammalian Cell Selection: | Puromycin  |
| Vector:                   | pLenti-C-Myc-DDK-P2A-Puro (PS100092)   |
| Tag:                      | Myc-DDK  |
| ACCN:                     | NM_031432  |
| ORF Size:                 | 831 bp   |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(RC220876).   |
| OTI Disclaimer:           | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a> |
| OTI Annotation:           | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| RefSeq:                   | <a href="#">NM_031432.1</a>  |
| RefSeq Size:              | 2160 bp  |
| RefSeq ORF:               | 834 bp   |
| Locus ID:                 | 83549  |
| UniProt ID:               | <a href="#">Q9HA47</a>   |
| Cytogenetics:             | 9q34.13  |
| Domains:                  | PRK  |
| Protein Families:         | Druggable Genome   |



[View online »](#)

**Protein Pathways:** Drug metabolism - other enzymes, Metabolic pathways, Pyrimidine metabolism

**MW:** 31.3 kDa

**Gene Summary:** This gene encodes a uridine-cytidine kinase that catalyzes the phosphorylation of uridine and cytidine to uridine monophosphate (UMP) and cytidine monophosphate (CMP) but not the phosphorylation of deoxyribonucleosides or purine ribonucleosides. This enzyme can also phosphorylate uridine and cytidine analogs and uses both ATP and GTP as a phosphate donor. Alternative splicing results in multiple splice variants encoding distinct isoforms. [provided by RefSeq, May 2012]