

Product datasheet for **RC220900**

KIAA1967 (CCAR2) (NM_021174) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	KIAA1967 (CCAR2) (NM_021174) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	KIAA1967
Synonyms:	DBC-1; DBC1; KIAA1967; NET35; p30 DBC; p30DBC
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>RC220900 representing NM_021174
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGTCCCAGTTTAAAGCGCCAGCGGATCAACCCGCTTCCAGGGGGACGCAACTTCTCAGGCACAGCTTCAA
 CATCTCTTCTGGGCCCTCCTCCTGGTTTGTCTACTCCTCCTGTGGCCACAGAAGTGTCCAGAATGCCAG
 GCACCTTCAGGGTGGGGAGAAACAGCGGTCTTACTGGTATTGTTACCAGCTTGCATGACTACTTTGGG
 GTTGTGGATGAAGAGGTCTTTTTTCAGCTAAGTGTGGTGAAGGGCCGTCTGCCCCAGCTGGGTGAGAAGG
 TGCTGGTGAAGGCTGCATAACAACCAGGCCAGGCAGTGCCTGGAATGCTGTCAAGGTGCAAACGCTCTC
 CAACCAGCCCCTACTGAAGTCCCAGCACCTCCTCTTCTGCATGTAGCAGCCCTGGGCCAGAAGCAAGGG
 ATCCTGGGAGCTCAGCCTCAGTTGATCTCCAGCCTCACCGATTCCCCACTCTTCTCAGAAGCCTC
 TGAGTCTCTTCCAACATCCCACACACTCACCTGAGCCACTGAACAGATTTCTGCCCGGGCCCTCA
 TGGACGGTTGGATCAGGGCCGAAGTGTACTATGACTCCAAGAAACGCAACAGCGGGCTGGTGGAGAG
 CCCTGGGGTGTAAAGCAAGGCATGACCTGCCTCCTTACCGGTCCACTCACTCTTACACTGTGG
 ACAGCCCCATCTGTGACTTCTAGAACTCCAGCGCCGTTACCGCAGCCTCCTGGTCCCCTCAGATTTTCT
 GTCCGTGCATCTGAGTTGGTATCAGCCTTCCCCTGAGCCAGCCCTTTTCCCTCCATCATCCAAGCCGG
 ATCCAGGTCTCTTCTGAAAAGGAGGCAGCTCCAGACGCTGGTGTGAGCCATCACTGCAGACAGTGACC
 CCGCTTATAGTTCGAAGTACTGCTGCTCTTCCCCGGGGTGGAGGAATTGTATCGTTGTTGCATGCT
 CTTTGTGGATGACATGGCTGAGCCAAGGGAGACGCCAGAGCATCCTCTGAAGCAGATTAAGTTTTGCTG
 GGCAGGAAAGAAGAGGAGGCAGTGTGGTGGGGTGAATGGTCTCCTTCCCTGGATGCCCTCGACCCCC
 AGGCTGACCCGAGGTGCTGGTGGTACCGCATCCGCTGTGCGCAGGCCAGACTGCCATTGATTTGAG
 CGGCTGTACCAAGTGGTGGCGCTTTGCCGAGTTTCAGTACTGACCGGGACCCCCCGCGGCTTCAG
 ACAGTGGTGGTGTACCTGCCGATGTCTGGACCATCATGCCTACTTTGGAGGAGTGGGAGCCCTGTGCC
 AGCAGAAAGTGCAGAGGCAGCTCCCCAACCCAGGAGGCACAAGGGGAAACGGAGCCTACTGAACAGGC
 ACCTGATGCCTTGGAGCAAGCAGCAGACTTCTAGACGGAACGCAGAACTCCAGAGGCCACCACACAG
 CAGGAAACGGACTGATCTCCAGAGGCCCTCCACCCCCCTAGAACCTGCTGCATCGCACGCCCTG
 GCTGTGTAACCTGTCCCTCCATGGGATGTGGAGGATCGGAGGCCAAAGGAAAGGATCTCTTTGAGGT
 GATGGTGTGCCGAGCTGTTTCTGGAGATGCTCCAGAGGGATTTGGCTATAGAGTTTATAAGATGCTA
 CTGAGCCTTCTGAAAAGTCTGTCTCCACCTGAACCTGAGAAGGAGGCGGCCAAGGAAGAAGCCA
 CCAAGGAGGAAGAAGCCATCAAAGAGGAGGTGGTCAAGGAGCCCAAGGATGAGGCACAGAATGAGGGCCC
 GGCTACAGAGTCAGAGGCCCGCTGAAGGAGGATGGGCTTTTGCCAAACCACTCTCTTCTGGGGGAGAG
 GAAGAAGAAAAACCCCGGGGCGAGGCTTCTGAGGACCTGTGTGAGATGGCCCTGGACCCAGAAGTGTGC
 TTCTGAGGGATGATGGAGAGGAGGATTTGAGGAGCAAGCTGGAGGATTCGGAGTCCGGTCCGTTGTC
 CTCAAACAGTCAGAGATGGAGTTCTTCACTTACAGGACATGCCCAAGGAGCTGGATCCCTCTGCTGTG
 CTCCCCTAGACTGTCTGCTTGTCTTTGTGTTCTTTGATGCCAAGTGGTGGCTACTTGCACCGGCGAG
 ACTTAGAGAGGATCCTCCTTACCCTTGGGATCCGGCTCAGTGCAGAGCAGGCCAAGCAGCTGGTCAAGCAG
 GGTGGTGACCCAGAACATCTGCCAGTACCGGACCTTCACTACAGCCGCCAGGAGGCTGGATGGTGGC
 CTTCGAGGAGGTGCTCTTCCGAAACCTGGACCTGCTGCCCCCTCTGGGAAAAGCACGAAGCCAGGTG
 CTGCCCCACAGAACACAAGCCTTGGTGTCCACAATGGCAGCCTGATTAACGTGGGGAGCCTGTGCA
 GCGCGCGGAGCAGCAGGACAGCGCCGGCTCTACCTAGAGAACAAGATCCACACACTGGAGCTGAAGCTG
 GAGGAGAGCCATAACCGTTTCTCAGCCACTGAAGTAACCAATAAGACGCTGGCGGAGAGATGCAGGAGC
 TGCGAGTCCGGCTGGCGGAGGCCAGGAGACCGCCGGACGGCGGAGCGACAGAAGGCCAGCTCCAGCG
 GCTGCTCAGGAGCTCCGAGGCGTCTGACCCCCCTGCAGCTGGAGATCCAGCGGGTGGTGGAAAAGGCT
 GACAGCTGGGTGGAGAAGGAGGACCGGCACCTAGCAAC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATTAAGTTTAA

Protein Sequence: >RC220900 representing NM_021174
 Red=Cloning site Green=Tags(s)

MSQFKRQRINPLPGRNFSGTASTSLLGPPPGLLTPPVATELSQNARHLQGGEKQRVFTGIVTSLHDYFG
 VVDEEVFFQLSVVKGRPLQLGEKVLVKAAYNPGQAVPWNVAVKVTLSNQPLLKSPAPPLHVAALGQKQG
 ILGAQPQLIFQPHRIPPLFPQKPLSLFQTSHTLHLSHLNRFPARGPHGRLDQGRSDDYDSKKRQRAGGE
 PWGAKKPRHDLPPYRVHLTPYTVDSPICDFLELQRRYRSLVPSDFLSVHLSWLSAFPLSQPFLSHHPSR
 IQVSSEKEAAPDAGAEPITADSDPAYSSKVLSSPGLLELYRCCMLFVDDMAEPRETPEHPLKQIKFLL
 GRKEEEAVLVGGEWSPSLDGLDPQADPQVLRVTAIRCAQAQTGIDLSGCTKWRRFAEFQYLQPGPPRRLQ
 TVVVYLPDVWTIMPTLEEWEALCQQKAAEAAPTQEAQGETEPTEQAPDALEQAADTSRRNAETPEATTQ
 QETDLDLPEAPPPLEPAVIARPGCVNLSLHGIVEDRRPKERISFEVMVLAELFLEMLQRDFGYRVYKML
 LSLPEKVVSPPEPEKEEAKEEAATKEEEAIKEEVVKEPKDEAQNEGPATESEAPLKEDGLLPKPLSSGGE
 EEEKPRGEASEDLCEMALDPELLLRDDGEEEFAGAKLEDSEVRSVSNQSEMEFSSLQDMPKELDPSAV
 LPLDCLLAFVFDANWCGYLHRRDLERILLTLGIRLSAEQAKQLVSRVVTQNICQYRSLQYSRQEGLDGG
 LPEEVLFGNLDLLPPPSTKPGAAPTEHKALVSHNGSLINVGSLLRQAEQQDSGRLYLENKIHTLELKL
 EESHNRFSATEVTNKTLAEMQELRVRLAEAEETARTAERQKSQLQRLQLQELRRRLTPLQLEIQRVVEKA
 DSWEKEEPPASN

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

SgfI-MluI

Cloning Scheme:



ACCN: NM_021174

ORF Size: 2769 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_021174.6](#)

RefSeq Size: 4031 bp

RefSeq ORF: 2772 bp

Locus ID: 57805


UniProt ID: [Q8N163](#)

Cytogenetics: 8p21.3


MW: 102.7 kDa

Gene Summary:


Core component of the DBIRD complex, a multiprotein complex that acts at the interface between core mRNP particles and RNA polymerase II (RNAPII) and integrates transcript elongation with the regulation of alternative splicing: the DBIRD complex affects local transcript elongation rates and alternative splicing of a large set of exons embedded in (A + T)-rich DNA regions. Inhibits SIRT1 deacetylase activity leading to increasing levels of p53/TP53 acetylation and p53-mediated apoptosis. Inhibits SUV39H1 methyltransferase activity. As part of a histone H3-specific methyltransferase complex may mediate ligand-dependent transcriptional activation by nuclear hormone receptors. Plays a critical role in maintaining genomic stability and cellular integrity following UV-induced genotoxic stress. Regulates the circadian expression of the core clock components NR1D1 and ARNTL/BMAL1. Enhances the transcriptional repressor activity of NR1D1 through stabilization of NR1D1 protein levels by preventing its ubiquitination and subsequent degradation (PubMed:18235501, PubMed:18235502, PubMed:19131338, PubMed:19218236, PubMed:22446626, PubMed:23352644, PubMed:23398316). Represses the ligand-dependent transcriptional activation function of ESR2 (PubMed:20074560). Acts as a regulator of PCK1 expression and gluconeogenesis by a mechanism that involves, at least in part, both NR1D1 and SIRT1 (PubMed:24415752). Negatively regulates the deacetylase activity of HDAC3 and can alter its subcellular localization (PubMed:21030595). Positively regulates the beta-catenin pathway (canonical Wnt signaling pathway) and is required for MCC-mediated repression of the beta-catenin pathway (PubMed:24824780). Represses ligand-dependent transcriptional activation function of NR1H2 and NR1H3 and inhibits the interaction of SIRT1 with NR1H3 (PubMed:25661920). Plays an important role in tumor suppression through p53/TP53 regulation; stabilizes p53/TP53 by affecting its interaction with ubiquitin ligase MDM2 (PubMed:25732823). Represses the transcriptional activator activity of BRCA1 (PubMed:20160719). Inhibits SIRT1 in a CHEK2 and PSEM3-dependent manner and inhibits the activity of CHEK2 in vitro (PubMed:25361978).[UniProtKB/Swiss-Prot Function]

Product images:

Circular map for RC220900



Western blot validation of overexpression lysate (Cat# [LY412049]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC220900 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified CCAR2 protein (Cat# [TP320900]). The protein was produced from HEK293T cells transfected with CCAR2 cDNA clone (Cat# RC220900) using MegaTran 2.0 (Cat# [TT210002]).