

Product datasheet for RC220918L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Alkyl DHAP synthase (AGPS) (NM 003659) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Alkyl DHAP synthase (AGPS) (NM_003659) Human Tagged ORF Clone Lentiviral Particle

Symbol: Alkyl DHAP synthase

Synonyms: ADAP-S; ADAS; ADHAPS; ADPS; ALDHPSY; RCDP3

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_003659 **ORF Size:** 1974 bp

ORF Nucleotide

OTI Disclaimer:

_.

Sequence:

The ORF insert of this clone is exactly the same as(RC220918).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 003659.1

RefSeq Size: 2074 bp
RefSeq ORF: 1977 bp
Locus ID: 8540

UniProt ID: O00116

Cytogenetics: 2q31.2

Domains: FAD-oxidase_C, FAD_binding_4

Protein Pathways: Ether lipid metabolism, Metabolic pathways





Alkyl DHAP synthase (AGPS) (NM_003659) Human Tagged ORF Clone Lentiviral Particle – RC220918L3V

MW: 72.91 kDa

Gene Summary:

This gene is a member of the FAD-binding oxidoreductase/transferase type 4 family. It encodes a protein that catalyzes the second step of ether lipid biosynthesis in which acyldihydroxyacetonephosphate (DHAP) is converted to alkyl-DHAP by the addition of a long chain alcohol and the removal of a long-chain acid anion. The protein is localized to the inner aspect of the peroxisomal membrane and requires FAD as a cofactor. Mutations in this gene have been associated with rhizomelic chondrodysplasia punctata, type 3 and Zellweger syndrome. [provided by RefSeq, Jul 2008]