

Protein Sequence: >RC220968 representing NM_001039671
Red=Cloning site Green=Tags(s)

MPGSASKRRIPVSQPGMADPHQLFDDTSSAQSRYGAQRAPGGLSYPAASPTPHAAFLADPVSNMAMAYG
 SSLAAQ GKELVDKNIDRFIPITKLKYYFAVDTMYVGRKGLLFFPYLHQDWEVQYQQDTPVAPRFDVNAP
 DLYIPAMAFITYVLVAGLALGTQDRFSPDLLGLQASSALAWLTLEVLAILLSLYLVTVNTDLTTIDL VAF
 LGYKYVGMIGGVL MGLLFGKIGIYYLVLGWCCVAIFVFMIRTLRLKILADAAAEGVPVARGARNQLRMYLTM
 AVAAAQPMLMYWLTFFHLVR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_001039671

ORF Size: 897 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001039671.3](#)

RefSeq Size: 1193 bp

RefSeq ORF: 900 bp

Locus ID: 90522


Cytogenetics: 19q13.2

Protein Families: Transmembrane

MW: 32.6 kDa

Gene Summary: Involved in the anterograde traffic pathway from the endoplasmic reticulum to the plasma membrane and the organization of the Golgi architecture (By similarity). Plays a key role in targeting to neuronal dendrites receptors such as HTR1A (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RC220968