

# Product datasheet for RC220992L4V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RTN3 (NM\_201428) Human Tagged ORF Clone Lentiviral Particle

### **Product data:**

**Product Type:** Lentiviral Particles

Product Name: RTN3 (NM 201428) Human Tagged ORF Clone Lentiviral Particle

Symbol: RTN3

Synonyms: ASYIP; HAP; NSPL2; NSPLII; RTN3-A1

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_201428 **ORF Size:** 3039 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC220992).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 201428.2</u>

 RefSeq Size:
 4919 bp

 RefSeq ORF:
 3042 bp

 Locus ID:
 10313

 UniProt ID:
 095197

 Cytogenetics:
 11q13.1

**Protein Families:** Transmembrane

MW: 110.7 kDa







### **Gene Summary:**

This gene belongs to the reticulon family of highly conserved genes that are preferentially expressed in neuroendocrine tissues. This family of proteins interact with, and modulate the activity of beta-amyloid converting enzyme 1 (BACE1), and the production of amyloid-beta. An increase in the expression of any reticulon protein substantially reduces the production of amyloid-beta, suggesting that reticulon proteins are negative modulators of BACE1 in cells. Alternatively spliced transcript variants encoding different isoforms have been found for this gene, and pseudogenes of this gene are located on chromosomes 4 and 12. [provided by RefSeq, May 2012]