

## Product datasheet for RC220995L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## APOBEC3A (NM\_145699) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** APOBEC3A (NM\_145699) Human Tagged ORF Clone Lentiviral Particle

Symbol: APOBEC3A

Synonyms: A3A; ARP3; bK150C2.1; PHRBN

Mammalian Cell

Selection:

**ORF Size:** 

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

597 bp

Tag: Myc-DDK

**ACCN:** NM\_145699

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC220995).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 145699.2, NP 663745.1

 RefSeq Size:
 1349 bp

 RefSeq ORF:
 600 bp

 Locus ID:
 200315

 UniProt ID:
 P31941

 Cytogenetics:
 22q13.1

**MW:** 22.8 kDa







## **Gene Summary:**

This gene is a member of the cytidine deaminase gene family. It is one of seven related genes or pseudogenes found in a cluster, thought to result from gene duplication, on chromosome 22. Members of the cluster encode proteins that are structurally and functionally related to the C to U RNA-editing cytidine deaminase APOBEC1. The protein encoded by this gene lacks the zinc binding activity of other family members. The protein plays a role in immunity, by restricting transmission of foreign DNA such as viruses. One mechanism of foreign DNA restriction is deamination of foreign double-stranded DNA cytidines to uridines, which leads to DNA degradation. However, other mechanisms are also thought to be involved, as antiviral effect is not dependent on deaminase activity. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2012]