

Product datasheet for **RC220997L4V**

Cytochrome P450 2C9 (CYP2C9) (NM_000771) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Cytochrome P450 2C9 (CYP2C9) (NM_000771) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Cytochrome P450 2C9
Synonyms:	CPC9; CYP2C; CYP2C10; CYPIIC9; P450IIC9
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_000771
ORF Size:	1470 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC220997).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_000771.2
RefSeq Size:	1835 bp
RefSeq ORF:	1473 bp
Locus ID:	1559
UniProt ID:	P11712
Cytogenetics:	10q23.33
Domains:	p450
Protein Families:	Druggable Genome, P450



[View online »](#)

Protein Pathways: Arachidonic acid metabolism, Drug metabolism - cytochrome P450, Linoleic acid metabolism, Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Retinol metabolism

MW: 55.4 kDa

Gene Summary: This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by rifampin. The enzyme is known to metabolize many xenobiotics, including phenytoin, tolbutamide, ibuprofen and S-warfarin. Studies identifying individuals who are poor metabolizers of phenytoin and tolbutamide suggest that this gene is polymorphic. The gene is located within a cluster of cytochrome P450 genes on chromosome 10q24. [provided by RefSeq, Jul 2008]