

## Product datasheet for **RC221051**

### **AKT3 (NM\_005465) Human Tagged ORF Clone**

#### **Product data:**

Product Type:	Expression Plasmids
Product Name:	AKT3 (NM_005465) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	AKT3
Synonyms:	MPPH; MPPH2; PKB-GAMMA; PKBG; PRKBG; RAC-gamma; RAC-PK-gamma; STK-2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

**ORF Nucleotide Sequence:**

>RC221051 representing NM\_005465  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGCATCGCC**

ATGAGCGATGTTACCATTGTGAAAGAAGGTTGGGTTCCAGAAGAGGGGAGAATATATAAAAACTGGAGGC  
 CAAGATACTTCCTTTGAAGACAGATGGCTCATTATAGGATATAAAGAGAAACCTCAAGATGTGGATTT  
 ACCTTATCCCCTCAACAACCTTTTCAGTGGCAAAATGCCAGTTAATGAAAAAGAACGACCAAGCCAAAC  
 ACATTTATAATCAGATGTCTCCAGTGGACTACTGTTATAGAGAGAACATTTATGTAGATACTCCAGAGG  
 AAAGGGAAGAATGGACAGAAGCTATCCAGGCTGTAGCAGACAGACTGCAGAGGCAAGAAGAGGAGAGAAT  
 GAATTGTAGTCCAACCTCACAAATTGATAATATAGGAGAGGAAGAGATGGATGCCTCTACAACCCATCAT  
 AAAAGAAAGACAATGAATGATTTTGACTATTTGAAACTACTAGGTAAGGCATTTTGGGAAAGTTATTT  
 TGGTTCGAGAGAAGGCAAGTGGAAAATACTATGCTATGAAGATTCTGAAGAAAGAAGTCATTATTGCAA  
 GGATGAAGTGGCACACACTCTAACTGAAAGCAGAGTATTAAGAACAAGTACATCCCTTTTAAACATCC  
 TTGAAATATTCTTCCAGACAAAAGACCGTTTGTGTTTTGTGATGGAATATGTTAATGGGGCGAGCTGT  
 TTTTCCATTTGTGCGAGAGAGCGGGTCTCTGAGGACCGCACACGTTTCTATGGTGCAGAAATTGTCTC  
 TGCTTGGACTATCTACATTCGGAAAGATTGTGTACCGTATCTCAAGTTGGAGAATCTAATGCTGGAC  
 AAAGATGGCCACATAAAAATTACAGATTTTGGACTTTGCAAAGAAGGGATCACAGATGCAGCCACCATGA  
 AGACATTCTGTGGCACTCCAGAATATCTGGCACCAGAGGTGTTAGAAGATAATGACTATGGCCGAGCAGT  
 AGACTGGTGGGCGCTAGGGTGTGATGTAATGATGTGTGGGAGGTTACCTTTCTACAACCAGGAC  
 CATGAGAACTTTTGAATTAATATTAATGGAAGACATTAATTTCTCGAACACTCTCTCAGATGCAA  
 AATCATTGCTTTTCAGGGCTCTTGATAAAGGATCCAAATAAACCGCTTGGTGGAGGACAGATGATGCAA  
 AGAAATTATGAGACACAGTTTCTCTCTGGAGTAAACTGGCAAGATGATATGATAAAAAGCTTGTACCT  
 CCTTTTAAACCTCAAGTAACATCTGAGACAGATACTAGATATTTTATGAGAATTTACAGCTCAGACTA  
 TTACAATAACACCACCTGAAAAATATGATGAGGATGGTATGGACTGCATGGACAATGAGAGGCGGCCGCA  
 TTTCCCTCAATTTTCTACTCTGCAAGTGGACGAGAA

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:**

>RC221051 representing NM\_005465  
 Red=Cloning site Green=Tags(s)

MSDVTIVKEGWVQKRGEYIKNWRPRYFLLKTDGSFIGYKEKPDVDLPYPLNNSVAKCQLMKTERPKPN  
 TFIIRCLQWTTVIERTFHVDTPEREWEATEAIQAVADRLQRQEEERMNCSPTSQIDNIGEEEMDASTTHH  
 KRKTMNDFDYLKLLGKTFGKVLVREKASGKYAMKILKKEVIAKDEVAHTL TESRVLNTRHPFLTS  
 LKYSFQTKDRLCFVMEYVNGGELFFHLSRERVF SEDRTRFYGAEIVSALDYLHSGKIVYRDLKLENMLD  
 KDGHKIDTDFGLCKEGITDAATMKTFCGTPEYLAPVLEDNDYGRAVDWWGLGVVYEMMCGRLPFYNQD  
 HEKLFELILMEDIKFPRTLSSDAKSLLSGLLIKDPNKRLGGPDDAKEIMRHSFFSGVNWQDVYDKLVP  
 PFKPQVTSETDRYFDEEFTAQITITPPEKYDEDMDCMDNERRPHFPQFSYSASGRE

**TR**TRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:**

Sgfl-MluI

**Cloning Scheme:**

**ACCN:**

NM\_005465

**ORF Size:**

1437 bp

**OTI Disclaimer:**

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at [custsupport@origene.com](mailto:custsupport@origene.com) or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:**

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:**


The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**


1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq:	<a href="#">NM_005465.7</a>
RefSeq Size:	3588 bp
RefSeq ORF:	1440 bp
Locus ID:	10000
UniProt ID:	<a href="#">Q9Y243</a>
Cytogenetics:	1q43-q44
Domains:	pkinase, S_TK_X, TyrKc, PH, S_TKc
Protein Families:	Druggable Genome, ES Cell Differentiation/IPS, Protein Kinase
Protein Pathways:	Acute myeloid leukemia, Adipocytokine signaling pathway, Apoptosis, B cell receptor signaling pathway, Chemokine signaling pathway, Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, ErbB signaling pathway, Fc epsilon RI signaling pathway, Fc gamma R-mediated phagocytosis, Focal adhesion, Glioma, Insulin signaling pathway, Jak-STAT signaling pathway, MAPK signaling pathway, Melanoma, mTOR signaling pathway, Neurotrophin signaling pathway, Non-small cell lung cancer, Pancreatic cancer, Pathways in cancer, Progesterone-mediated oocyte maturation, Prostate cancer, Renal cell carcinoma, Small cell lung cancer, T cell receptor signaling pathway, Tight junction, Toll-like receptor signaling pathway, VEGF signaling pathway
MW:	55.6 kDa
Gene Summary:	The protein encoded by this gene is a member of the AKT, also called PKB, serine/threonine protein kinase family. AKT kinases are known to be regulators of cell signaling in response to insulin and growth factors. They are involved in a wide variety of biological processes including cell proliferation, differentiation, apoptosis, tumorigenesis, as well as glycogen synthesis and glucose uptake. This kinase has been shown to be stimulated by platelet-derived growth factor (PDGF), insulin, and insulin-like growth factor 1 (IGF1). Alternatively splice transcript variants encoding distinct isoforms have been described. [provided by RefSeq, Jul 2008]


Product images:




Circular map for RC221051




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY AKT3 (Cat# RC221051, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-AKT3 (Cat# [TA500001]).



Coomassie blue staining of purified AKT3 protein (Cat# [TP321051]). The protein was produced from HEK293T cells transfected with AKT3 cDNA clone (Cat# RC221051) using MegaTran 2.0 (Cat# [TT210002]).



Anti-AKT3 mouse monoclonal antibody ([TA500001]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY AKT3 (RC221051).



HEK293T cells transfected with either overexpress plasmid (RC221051, Red) or empty vector control plasmid (Blue) were immunostained by anti-AKT3 antibody ([TA500001]), and then analyzed by flow cytometry.