

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC221072L4V

ZFAND5 (NM_001102420) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	ZFAND5 (NM_001102420) Human Tagged ORF Clone Lentiviral Particle
Symbol:	ZFAND5
Synonyms:	ZA20D2; ZFAND5A; ZNF216
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001102420
ORF Size:	639 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC221072).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001102420.1, NP 001095890.1</u>
RefSeq Size:	5887 bp
RefSeq ORF:	642 bp
Locus ID:	7763
UniProt ID:	<u>076080</u>
Cytogenetics:	9q21.13
MW:	23.1 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Involved in protein degradation via the ubiquitin-proteasome system. May act by anchoring
ubiquitinated proteins to the proteasome. Plays a role in ubiquitin-mediated protein
degradation during muscle atrophy. Plays a role in the regulation of NF-kappa-B activation
and apoptosis. Inhibits NF-kappa-B activation triggered by overexpression of RIPK1 and
TRAF6 but not of RELA. Inhibits also tumor necrosis factor (TNF), IL-1 and TLR4-induced NF-
kappa-B activation in a dose-dependent manner. Overexpression sensitizes cells to TNF-
induced apoptosis. Is a potent inhibitory factor for osteoclast differentiation.
[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US