OriGene Technologies, Inc.
9620 Medical Center Drive, Ste 200
Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Product datasheet for RC221087L4V

EAAT4 (SLC1A6) (NM_005071) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	EAAT4 (SLC1A6) (NM_005071) Human Tagged ORF Clone Lentiviral Particle
Symbol:	EAAT4
Synonyms:	EAAT4
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_005071
ORF Size:	1692 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC221087).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_005071.1
RefSeq Size:	1719 bp
RefSeq ORF:	1695 bp
Locus ID:	6511
UniProt ID:	P48664
Cytogenetics:	19p13.12
Domains:	SDF
Protein Families:	Transmembrane

View online »

MW: 61.4 kDa

Gene Summary: Sodium-dependent, high-affinity amino acid transporter that mediates the uptake of L-glutamate and also L-aspartate and D-aspartate (PubMed:7791878). Functions as a symporter that transports one amino acid molecule together with two or three Na(+) ions and one proton, in parallel with the counter-transport of one K(+) ion. Mediates Cl(-) flux that is not coupled to amino acid transport; this avoids the accumulation of negative charges due to aspartate and Na(+) symport (By similarity). Plays a redundant role in the rapid removal of released glutamate from the synaptic cleft, which is essential for terminating the postsynaptic action of glutamate (Probable).[UniProtKB/Swiss-Prot Function]