


# Product datasheet for RC221108L1

# SMN1 (NM\_022874) Human Tagged Lenti ORF Clone

## **Product data:**

#### **Product Type: Expression Plasmids Product Name:** SMN1 (NM 022874) Human Tagged Lenti ORF Clone Tag: Myc-DDK Symbol: SMN1 Synonyms: BCD541; GEMIN1; SMA; SMA1; SMA2; SMA3; SMA4; SMA@; SMN; SMNT; T-BCD541; TDRD16A Mammalian Cell None Selection: Vector: pLenti-C-Myc-DDK (PS100064) E. coli Selection: Chloramphenicol (34 ug/mL) The ORF insert of this clone is exactly the same as(RC221108). **ORF** Nucleotide Sequence: **Restriction Sites:** Sgfl-Mlul **Cloning Scheme:** Cloning sites used for ORF Shuttling: ORF Sqf I Mlu I --- GCG ATC GC C ATG --- //--- NNN ACG CGT ---



ACCN: ORF Size: NM\_022874 786 bp

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



View online »

1/4


	(NM_022874) Human Tagged Lenti ORF Clone – RC221108L1
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
RefSeq:	<u>NM 022874.1</u>
RefSeq Size:	1525 bp
RefSeq ORF:	789 bp
Locus ID:	6606
UniProt ID:	<u>Q16637</u>
Cytogenetics:	5q13.2
Domains:	TUDOR
Protein Families:	Druggable Genome, Stem cell - Pluripotency
MW:	28.4 kDa

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### SMN1 (NM\_022874) Human Tagged Lenti ORF Clone – RC221108L1

This gene is part of a 500 kb inverted duplication on chromosome 5q13. This duplicated Gene Summary: region contains at least four genes and repetitive elements which make it prone to rearrangements and deletions. The repetitiveness and complexity of the sequence have also caused difficulty in determining the organization of this genomic region. The telomeric and centromeric copies of this gene are nearly identical and encode the same protein. However, mutations in this gene, the telomeric copy, are associated with spinal muscular atrophy; mutations in the centromeric copy do not lead to disease. The centromeric copy may be a modifier of disease caused by mutation in the telomeric copy. The critical sequence difference between the two genes is a single nucleotide in exon 7, which is thought to be an exon splice enhancer. Note that the nine exons of both the telomeric and centromeric copies are designated historically as exon 1, 2a, 2b, and 3-8. It is thought that gene conversion events may involve the two genes, leading to varying copy numbers of each gene. The protein encoded by this gene localizes to both the cytoplasm and the nucleus. Within the nucleus, the protein localizes to subnuclear bodies called gems which are found near coiled bodies containing high concentrations of small ribonucleoproteins (snRNPs). This protein forms heteromeric complexes with proteins such as SIP1 and GEMIN4, and also interacts with several proteins known to be involved in the biogenesis of snRNPs, such as hnRNP U protein and the small nucleolar RNA binding protein. Multiple transcript variants encoding distinct isoforms have been described. [provided by RefSeq, Jul 2014]


### **Product images:**



Circular map for RC221108L1

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US





Double digestion of RC221108L1 using Sgfl and Mlul

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US