

Product datasheet for **RC221360L3V**

MRC2 (NM_006039) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	MRC2 (NM_006039) Human Tagged ORF Clone Lentiviral Particle
Symbol:	MRC2
Synonyms:	CD280; CLEC13E; ENDO180; UPARAP
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_006039
ORF Size:	4437 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC221360).
OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info</p>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_006039.2 , NP_006030.1
RefSeq Size:	4700 bp
RefSeq ORF:	4440 bp



[View online »](#)

Locus ID:	9902
UniProt ID:	Q9UBG0
Cytogenetics:	17q23.2
Protein Families:	Druggable Genome, Transmembrane
MW:	166.5 kDa
Gene Summary:	<p>This gene encodes a member of the mannose receptor family of proteins that contain a fibronectin type II domain and multiple C-type lectin-like domains. The encoded protein plays a role in extracellular matrix remodeling by mediating the internalization and lysosomal degradation of collagen ligands. Expression of this gene may play a role in the tumorigenesis and metastasis of several malignancies including breast cancer, gliomas and metastatic bone disease. [provided by RefSeq, Feb 2012]</p>