

## Product datasheet for RC221369L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## UCP4 (SLC25A27) (NM 004277) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type: Lentiviral Particles** 

**Product Name:** UCP4 (SLC25A27) (NM 004277) Human Tagged ORF Clone Lentiviral Particle

Symbol: UCP4 Synonyms: **Mammalian Cell** 

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag: NM 004277 ACCN:

**ORF Size:** 969 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC221369).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 004277.3, NP 004268.3

RefSeq Size: 2959 bp RefSeq ORF: 972 bp Locus ID: 9481 **UniProt ID:** O95847 Cytogenetics: 6p12.3

**Domains:** mito\_carr

**Protein Families:** Druggable Genome





MW: 35.9 kDa

**Gene Summary:** 

Mitochondrial uncoupling proteins (UCP) are members of the larger family of mitochondrial anion carrier proteins (MACP). UCPs separate oxidative phosphorylation from ATP synthesis with energy dissipated as heat, also referred to as the mitochondrial proton leak. UCPs facilitate the transfer of anions from the inner to the outer mitochondrial membrane and the return transfer of protons from the outer to the inner mitochondrial membrane. They also reduce the mitochondrial membrane potential in mammalian cells. Tissue specificity occurs for the different UCPs and the exact methods of how UCPs transfer H+/OH- are not known. UCPs contain the three homologous protein domains of MACPs. Transcripts of this gene are only detected in brain tissue and are specifically modulated by various environmental conditions. Alternative splicing results in multiple transcript variants.[provided by RefSeq, Feb 2011]