

Product datasheet for RC221381L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Caspase-6 (CASP6) (NM_032992) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Caspase-6 (CASP6) (NM 032992) Human Tagged ORF Clone Lentiviral Particle

Symbol: Caspase-6

Synonyms: MCH2

Mammalian Cell None

Selection:

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_032992

ORF Size: 612 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC221381).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 032992.2

 RefSeq Size:
 1394 bp

 RefSeq ORF:
 615 bp

 Locus ID:
 839

 UniProt ID:
 P55212

Cytogenetics: 4q25

Protein Families: Druggable Genome, Protease, Stem cell - Pluripotency

Protein Pathways: Apoptosis





MW: 22.4 kDa

Gene Summary:

This gene encodes a member of the cysteine-aspartic acid protease (caspase) family of enzymes. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic acid residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein is processed by caspases 7, 8 and 10, and is thought to function as a downstream enzyme in the caspase activation cascade. Alternative splicing of this gene results in multiple transcript variants that encode different isoforms. [provided by RefSeq, Oct 2015]