

Product datasheet for RC221399L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

UBF1 (UBTF) (NM_014233) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: UBF1 (UBTF) (NM_014233) Human Tagged ORF Clone Lentiviral Particle

Symbol: UBF1

Synonyms: CONDBA; NOR-90; UBF; UBF-1; UBF1; UBF2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_014233 **ORF Size:** 2292 bp

ORF Nucleotide

2232 00

Sequence:

The ORF insert of this clone is exactly the same as(RC221399).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 014233.1

 RefSeq Size:
 3097 bp

 RefSeq ORF:
 2295 bp

 Locus ID:
 7343

 UniProt ID:
 P17480

 Cytogenetics:
 17q21.31

Domains: HMG

Protein Families: Transcription Factors





ORIGENE

MW: 89.2 kDa

Gene Summary:

This gene encodes a member of the HMG-box DNA-binding protein family. The encoded protein plays a critical role in ribosomal RNA transcription as a key component of the preinitiation complex, mediating the recruitment of RNA polymerase I to rDNA promoter regions. The encoded protein may also play important roles in chromatin remodeling and pre-rRNA processing, and its activity is regulated by both phosphorylation and acetylation. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. Pseudogenes of this gene are located on the short arm of chromosomes 3, 11 and X and the long arm of chromosome 11. [provided by RefSeq, Aug 2011]