

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC221493L3V

## Caspase-7 (CASP7) (NM\_033340) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Caspase-7 (CASP7) (NM_033340) Human Tagged ORF Clone Lentiviral Particle
Symbol:	CASP7
Synonyms:	CASP-7; CMH-1; ICE-LAP3; LICE2; MCH3
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_033340
ORF Size:	759 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC221493).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 033340.2</u>
RefSeq Size:	2433 bp
RefSeq ORF:	762 bp
Locus ID:	840
UniProt ID:	<u>P55210</u>
Cytogenetics:	10q25.3
Domains:	Peptidase_C14
Protein Families:	Druggable Genome, Protease



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

<b>GRIGENE</b> Caspase-7 (CASP7) (NM_033340) Human Tagged ORF Clone Lentiviral Particle – RC221493L3V	
Protein Pathways:	Alzheimer's disease, Apoptosis
MW:	27.8 kDa
Gene Summary:	This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. The precursor of the encoded protein is cleaved by caspase 3 and 10, is activated upon cell death stimuli and induces apoptosis. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, May 2012]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US