

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC221498L2V

## Protein Kinase D2 (PRKD2) (NM\_001079881) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Protein Kinase D2 (PRKD2) (NM_001079881) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Protein Kinase D2
Synonyms:	HSPC187; nPKC-D2; PKD2
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_001079881
ORF Size:	2634 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC221498).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001079881.1, NP 001073350.1</u>
RefSeq Size:	3202 bp
RefSeq ORF:	2637 bp
Locus ID:	25865
UniProt ID:	<u>Q9BZL6</u>
Cytogenetics:	19q13.32
Protein Families:	Druggable Genome, Protein Kinase
MW:	96.5 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:The protein encoded by this gene belongs to the protein kinase D (PKD) family of<br/>serine/threonine protein kinases. This kinase can be activated by phorbol esters as well as by<br/>gastrin via the cholecystokinin B receptor (CCKBR) in gastric cancer cells. It can bind to<br/>diacylglycerol (DAG) in the trans-Golgi network (TGN) and may regulate basolateral<br/>membrane protein exit from TGN. Alternative splicing results in multiple transcript variants<br/>encoding different isoforms. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US