

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC221534L4V

C18orf1 (LDLRAD4) (NM_001003674) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	C18orf1 (LDLRAD4) (NM_001003674) Human Tagged ORF Clone Lentiviral Particle
Symbol:	C18orf1
Synonyms:	C18orf1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001003674
ORF Size:	807 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC221534).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001003674.1, NP 001003674.1</u>
RefSeq Size:	8944 bp
RefSeq ORF:	810 bp
Locus ID:	753
UniProt ID:	<u>O15165</u>
Cytogenetics:	18p11.21
Protein Families:	Druggable Genome, Transmembrane
MW:	29.9 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Functions as a negative regulator of TGF-beta signaling and thereby probably plays a role in
cell proliferation, differentiation, apoptosis, motility, extracellular matrix production and
immunosuppression. In the canonical TGF-beta pathway, ZFYVE9/SARA recruits the
intracellular signal transducer and transcriptional modulators SMAD2 and SMAD3 to the TGF-
beta receptor. Phosphorylated by the receptor, SMAD2 and SMAD3 then form a heteromeric
complex with SMAD4 that translocates to the nucleus to regulate transcription. Through
interaction with SMAD2 and SMAD3, LDLRAD4 may compete with ZFYVE9 and SMAD4 and
prevent propagation of the intracellular signal.[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US