



OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_003048.3
RefSeq Size:	5446 bp
RefSeq ORF:	2439 bp
Locus ID:	6549
UniProt ID:	Q9UBY0
Cytogenetics:	2q12.1
Domains:	Na_H_Exchanger
Protein Families:	Druggable Genome, Transmembrane
MW:	91.3 kDa
Gene Summary:	This gene encodes a member of the sodium-hydrogen exchanger (NHE) protein family. These proteins are involved in sodium-ion transport by exchanging intracellular hydrogen ions to external sodium ions and help in the regulation of cell pH and volume. The encoded protein is localized to the apical membrane and is involved in apical absorption of sodium. [provided by RefSeq, Jun 2016]

Product images:



Circular map for RC221550L1



Double digestion of RC221550L1 using SgfI and MluI