

## Product datasheet for RC221589L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Caspase-7 (CASP7) (NM 033338) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: Caspase-7 (CASP7) (NM\_033338) Human Tagged ORF Clone Lentiviral Particle

Symbol: Caspase-7

Synonyms: CASP-7; CMH-1; ICE-LAP3; LICE2; MCH3

Mammalian Cell

Selection:

ACCN:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 033338

Tag: Myc-DDK

ORF Size: 1008 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC221589).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 033338.4

RefSeq Size: 2712 bp
RefSeq ORF: 1011 bp

 Locus ID:
 840

 UniProt ID:
 P55210

Cytogenetics: 10q25.3

**Protein Families:** Druggable Genome, Protease

**Protein Pathways:** Alzheimer's disease, Apoptosis





**MW:** 37.6 kDa

**Gene Summary:** 

This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. The precursor of the encoded protein is cleaved by caspase 3 and 10, is activated upon cell death stimuli and induces apoptosis. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, May 2012]