

## Product datasheet for RC221599L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CD1 (CD1A) (NM\_001763) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: CD1 (CD1A) (NM\_001763) Human Tagged ORF Clone Lentiviral Particle

Symbol: CD<sup>2</sup>

Synonyms: CD1; FCB6; HTA1; R4; T6

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 001763

ORF Size: 981 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC221599).

Sequence:

Cytogenetics:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001763.1, NP 001754.1

 RefSeq Size:
 2072 bp

 RefSeq ORF:
 984 bp

 Locus ID:
 909

 UniProt ID:
 P06126

**Protein Families:** Druggable Genome, Transmembrane

1q23.1

**Protein Pathways:** Hematopoietic cell lineage







**MW:** 37.17 kDa

**Gene Summary:** 

This gene encodes a member of the CD1 family of transmembrane glycoproteins, which are structurally related to the major histocompatibility complex (MHC) proteins and form heterodimers with beta-2-microglobulin. The CD1 proteins mediate the presentation of primarily lipid and glycolipid antigens of self or microbial origin to T cells. The human genome contains five CD1 family genes organized in a cluster on chromosome 1. The CD1 family members are thought to differ in their cellular localization and specificity for particular lipid ligands. The protein encoded by this gene localizes to the plasma membrane and to recycling vesicles of the early endocytic system. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016]